GUAIFENESIN AND CODEINE PHOSPHATE- guaifenesin and codeine phosphate solution QUAGEN PHARMACEUTICALS LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Guaifenesin and Codeine Phosphate Oral Solution, USP CV 100 mg/10 mg per 5 mL

SUGAR FREE, DYE FREE, ALCOHOL FREE

Drug Facts

Active ingredients		
(in each $5 \text{ mL} = 1 \text{ tsp}$)	Purpose	
Codeine phosphate, USP 10 mg		
	Cough suppressant	
Guaifenesin, USP 100 mg		
3	Expectorant	

Uses

- temporarily relieves:
- cough due to minor throat and bronchial irritation as may occur with a cold or inhaled irritants
- your cough to help you sleep
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

Warnings

Ask your doctor before use if

- you have a persistent cough, this may be a sign of a serious condition
- you have a persistent cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- you have a cough that is accompanied by excessive phlegm (mucus)
- you have chronic pulmonary disease or shortness of breath
- giving to a child who is taking other drugs

When using this product

- giving a higher dose than recommended by a doctor could result in serious side effects for your child. A special measuring device should be used to give an accurate dose of this product to children under 6 years of age.
- may cause or aggravate constipation

Stop use and ask a doctor if

• symptoms do not improve within 7 days, tend to recur or are accompanied by fever and rash or persistent headache. These may be symptoms of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Do not exceed 6 doses in 24 hours.

Adults and children 12 years of age and	2 tsp (10 mL) every 4 hours,
over:	or as directed by a doctor.
Children 6 to under 12 years of age:	1 tsp (5 mL) every 4 hours,
	or as directed by a doctor.
Children under 6 years of age:	Consult a doctor.

Other information

Each tsp (5 mL) contains 3 mg sodium.

Store at controlled room temperature 15°-30°C (59°-86°F). You may report side effects by calling 1-888-344-9603 or FDA at 1-800-FDA-1088.

Inactive ingredients

Caramel flavor, cherry flavor, citric acid, glycerin, peppermint flavor, purified water, sodium benzoate, sodium saccharin, sorbitol.

Tamper evident by foil seal under cap. Do not use if foil seal is broken or missing.

Manufactured by:

Quagen Pharmaceuticals LLC West Caldwell, NJ 07006

MADE IN USA

PRINCIPAL DISPLAY PANEL-4 fl oz (118 mL) Container Label



ZM

Purpose

Tamper evident by foil seal under cap

MADE IN USA

PEEL HERE

100 mg/10 mg per 5 mL

Phosphate Oral Solution, USP 🚺

Expectorant/Cough Suppressant

SUGAR FREE, DYE FREE, ALCOHOL FREE

Each 5 mL (1 teaspoonful) contains:

Codeine phosphate, USP...... 10 mg Guaifenesin, USP 100 mg

(WARNING: May be habit-forming)

4 fl oz (118 mL)

Drug Facts

Active ingredients

(in each 5 mL= 1 tsp)

Codeine phosphate, USP 10 mg

Guaifenesin, USP 100 mg....

Uses temporarily relieves:

- cough due to minor throat and bronchial irritation as may occur with a cold or inhaled irritants
- your cough to help you sleep
- in helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.

Warnings

Ask your doctor before use if

- you have a persistent cough, this may be a sign of a serious condition
- you have a persistent cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
 you have a cough that is accompanied by excessive phlegm (mucus)
- you have chronic pulmonary disease or shortness of breath
- giving to a child who is taking other drugs

When using this product

- qiving a higher dose than recommended by a doctor could result in serious side effects for your child. A special measuring device should be use to give an accurate dose of this product children under 6 years of age.
- may cause or aggravate constipation

symptoms do not improve within 7 days, tend to recur or are accompanied by fever and rash or persistent headache. These may be symptoms of a serious condition.

Keep out of the reach of children, in case of overdose, get medical help or contact a Poison Control Center right away.

Directions

Do not exceed 6 doses in 24 hours.

Adults and children 12 years of age and over: 2 tsp (10 mL) every 4 hours, or as directed by a doctor. Children 6 to under 12 years of age: 1 tsp (5 mL) every 4 hours, or as directed by a doctor. Children under 6 years of age: Consult a doctor.

Other information

Each tsp (5 mL) contains 3 mg sodium.

Store at controlled room temperature 15°-30°C (59°-86°F). You may report side effects by calling 1-888-344-9603 or FDA at 1-800-FDA-1088.

Inactive ingredients

Caramel flavor, cherry flavor, citric acid, glycerin, peppermint flavor, purified water, sodium benzoate, sodium saccharin, sorbitol

Manufactured by:

Quagen Pharmaceuticals LLC West Caldwell, NJ 07006

Drug Facts (continued) Stop use and ask a doctor if

If pregnant or breast-feeding, ask a health professional before use.

PRINCIPAL DISPLAY PANEL-8 fl oz (237 mL) Container Label



ALCOHOL FREE

Each 5 mL (1 teaspoonful) contains: Codeine phosphate, USP...... 10 mg Guaifenesin, USP 100 mg

(WARNING: May be habit-forming) Tamper evident by foil seal under cap. Do not use if foil seal is broken or missing.

8 fl oz (237 mL)

Drug Facts

Active ingredients

(in each 5 mL = 1 tsp)

Codeine phosphate, USP 10 mg...... Cough suppressant Guaifenesin, USP 100 mg

Uses ■ temporarily relieves:

- cough due to minor throat and bronchial irritation as may occur with a cold or inhaled irritants
- your cough to help you sleep
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Ask your doctor before use if

- you have a persistent cough, this may be a sign of a serious condition
- you have a persistent cough such as occurs with smoking, asthma, chronic bronchitis or emphysema you have a cough that is accompanied by excessive
- phlegm (mucus) you have chronic pulmonary disease or shortness of breath giving to a child who is taking other drugs

When using this product

- giving a higher dose than recommended by a doctor could result in serious side effects for your child. A special measuring device should be used to give an accurate dose of this product to children under 6 years of age.
- may cause or aggravate constination

Drug Facts (continued)

Stop use and ask a doctor if

symptoms do not improve within 7 days, tend to recur or are accompanied by fever and rash or persistent headache. These may be symptoms of a serious condition.

f pregnant or breast-feeding, ask a health professional before use Keep out of the reach of children, in case of overdose, get medical help or contact a Poison Control Center right away.

Purpose

Directions
■ Do not exceed 6 doses in 24 hours.

Adults and children 12 2 tsp (10 mL) every 4 hours. years of age and over: Children 6 to under 12 or as directed by a doctor. 1 tsp (5 mL) every 4 hours. ears of age or as directed by a doctor. Children under 6 years of age: | Consult a doctor

Other information

Each tsp (5 mL) contains 3 mg sodium. Store at controlled room temperature 15°-30°C (59°-86°F). You may report side effects by calling 1-888-344-9603 or FDA at 1-800-FDA-1088.

Inactive ingredients

Caramel flavor, cherry flavor, citric acid, glycerin, peppermint flavor, purified water, sodium benzoate, sodium saccharin, sorbitol.

Manufactured by: Quagen Pharmaceuticals LLC

West Caldwell, NJ 07006



50097 Rev. 06/21



0.75" X 3.0" NON VARNISH AREA

PRINCIPAL DISPLAY PANEL-16 fl oz (473 mL) Container Label



NDC 70752-180-12

Guaifenesin and **Codeine Phosphate** Oral Solution, USP (V

100 mg/10 mg per 5 mL

Expectorant/Cough Suppressant

SUGAR FREE, DYE FREE, **ALCOHOL FREE**

Each 5 mL (1 teaspoonful) contains: Codeine phosphate, USP...... 10 mg

Guaifenesin, USP 100 mg (WARNING: May be habit-forming)

Tamper evident by foil seal under cap. Do not use if foil seal is broken or missing.

16 fl oz (473 mL)

Drug Facts

Active ingredients

(in each 5 mL= 1 tsp) Codeine phosphate, USP 10 mg Cough suppressant Guaifenesin, USP 100 mg....

Purpose

Uses ■ temporarily relieves:

- cough due to minor throat and bronchial irritation as may occur with a cold or inhaled irritants
- your cough to help you sleep
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more

Warnings

Ask your doctor before use if

- you have a persistent cough, this may be a sign of a serious condition
- you have a persistent cough such as occurs with
- smoking, asthma, chronic bronchitis or emphysema you have a cough that is accompanied by
- excessive phleam (mucus) you have chronic pulmonary disease or shortness of breath
- giving to a child who is taking other drugs

When using this product

 giving a higher dose than recommended by a doctor could result in serious side effects for your child. A special measuring device should be used to give an accurate dose of this product to children under 6 years of age.

may cause or aggravate constipation

Drug Facts (continued)

Stop use and ask a doctor if

symptoms do not improve within 7 days, tend to recur or are accompanied by fever and rash or persistent headache. These may be symptoms of a serious condition.

If pregnant or breast-feeding, ask a health professional

Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Do not exceed 6 doses in 24 hours
Adults and children 12 2 tsp 2 tsp (10 mL) every 4 hours, ears of age and over or as directed by a doctor. Chi**l**dren 6 to under 12 1 tsp (5 mL) every 4 hours, or as directed by a doctor. vears of age: Children under 6 years of age: Consult a doctor.

Other information

Each tsp (5 mL) contains 3 mg sodium Store at controlled room temperature 15°-30°C 59°-86°F). You may report side effects by calling 1-888-344-9603 or FDA at 1-800-FDA-1088.

Inactive ingredients

Caramel flavor, cherry flavor, citric acid, glycerin, peppermint flavor, purified water, sodium benzoate, sodium saccharin, sorbitol,

Manufactured by: Quagen Pharmaceuticals LLC West Caldwell, NJ 07006



50099



10" X 40" NON VARNISH AREA

NDC:70752-180

GUAIFENESIN AND CODEINE PHOSPHATE

guaifenesin and codeine phosphate solution

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg in 5 mL
CODEINE PHOSPHATE (UNII: GSL05Y1MN6) (CODEINE ANHYDROUS - UNII: UX6OWY2V7J)	CODEINE PHOSPHATE	10 mg in 5 mL

Inactive Ingredients			
Ingredient Name	Strength		
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)			
GLYCERIN (UNII: PDC6A3C0OX)			
WATER (UNII: 059QF0KO0R)			
SODIUM BENZOATE (UNII: OJ245FE5EU)			
SACCHARIN SODIUM (UNII: SB8ZUX40TY)			
SORBITOL (UNII: 506T60A25R)			

Product Characteristics		
Color		Score
Shape		Size
Flavor	CARAMEL, CHERRY, PEPPERMINT	Imprint Code
Contains		

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70752-180- 06	118 mL in 1 BOTTLE; Type 0: Not a Combination Product	05/23/2022	
2	NDC:70752-180- 15	237 mL in 1 BOTTLE; Type 0: Not a Combination Product	05/23/2022	
3	NDC:70752-180- 12	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	05/23/2022	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Marketing E Date Date	
OTC monograph final	part341	05/23/2022	

Labeler - QUAGEN PHARMACEUTICALS LLC (073645339)

Establishment				
Name	Address	ID/FEI	Business Operations	
QUAGEN PHARMACEUTICALS LLC		080281331	manufacture(70752-180) , pack(70752-180)	

Revised: 7/2022 QUAGEN PHARMACEUTICALS LLC